

Spectral Analysis of Super Soft X-ray Sources: V4743 Sagittarii

T. Rauch,^{1,2} M. Orio,^{3,4} R. Gonzales-Riestra,⁵ and M. Still⁶

¹*Dr.-Remeis-Sternwarte, Sternwartstraße 7, 96049 Bamberg, Germany*

²*Institut für Astronomie und Astrophysik, Sand 1, 72076 Tübingen, Germany*

³*Istituto Nazionale di Astrofisica (INAF), Osservatorio Astronomico di Torino, Strada Osservatorio, 20, 10025 Pino Torinese (TO), Italy*

⁴*Department of Astronomy, 475 N. Charter Street, University of Wisconsin, Madison, WI 53706, USA*

⁵*XMM-Newton Science Operations Centre, Villafranca del Castillo, P.O. Box - Apdo. 50727, 28080 Madrid, Spain*

⁶*NASA/GSFC, Code 662, Greenbelt, MD 20771, USA*

Abstract. Half a year after its outburst, the nova V4743 Sgr evolved into the brightest super-soft X-ray source in the sky with a flux maximum around 30 Å, exhibiting resonance lines of C V, C VI, N VI, N VII, and O VII. We present preliminary results of an analysis of the XMM-Newton RGS spectra by means of NLTE model-atmosphere techniques.

1. Introduction

Classical novae occur in close binary systems (main-sequence star + white dwarf). V4743 Sgr has been discovered as a nova on Sep 20, 2002 (Haseda et al. 2002). A first CHANDRA observation, taken with ACIS-S in Dec 2002 (Ness et al. 2003), shows a X-ray spectrum which is consistent with a view on the optically thick atmosphere of an extremely hot white dwarf with strong C V, N VI, and N VII absorption features (indicating T_{eff} of 1 – 2 MK, Ness et al. 2003) which are blue shifted (-2400 km/sec) — most likely expanding gas which consists out of CNO-processed material.

2. Observations

V4743 Sgr was a very fast nova, with a steep decline in the optical light curve and large ejection velocity. The time to decay by 3 mag in the visual (t_3), was 15 days and the Full Width at Half Maximum (FWHM) of the H α line reached 2400 km/sec (Kato et al. 2002).

V4743 Sgr was observed with the XMM-Newton satellite on Apr 4, 2003 for about 10 h. The EPIC-pn count rate in timing mode was 1348 cts/sec (Orio et al. 2003). The spectrum appears to be extremely soft with absorption features (Fig. 3), which are attributed to the white dwarf's atmosphere. Because of the high S/N achieved, we have used the XMM-Newton RGS spectra obtained




Figure 1. Synthetic fluxes calculated from line-blanketed NLTE model atmospheres including H, He, C, N, and O. The abundances are mass ratios.

during this observation to start fitting the stellar atmosphere, although we are also working at fitting the previous spectrum, and later ones, obtained with CHANDRA LETG.

3. NLTE Model Atmospheres

The plane-parallel, static models are calculated with TMAP, the Tübingen NLTE Model Atmosphere Package (Werner et al. 2003, Rauch & Deetjen 2003). The first set of models is composed out of H+He+C+N+O, then Ne+Mg+S are added subsequently in a line-formation calculation. However, in general




Figure 2. Comparison of fluxes of $T_{\text{eff}} = 700 \text{ kK}$ models with different abundance ratios (thick: like given in Fig. 1, thin: solar, dashed: black body).

Table 1. Statistics of the model atoms used in our calculations of the NLTE model atmospheres. The notation is: NLTE = levels treated in NLTE, LTE = LTE levels, RBB = radiative bound-bound transitions

atom	ion	NLTE	LTE	RBB	atom	ion	NLTE	LTE	RBB
H	I	5	11	10	Ne	VII	10	50	12
	II	1	-	-		VIII	8	18	15
He	I	1	21	0		IX	5	6	3
	II	10	22	45		X	1	0	0
	III	1	-	-	Mg	VIII	1	19	0
C	V	29	21	60		IX	3	23	1
	VI	15	21	26		X	2	3	1
	VII	1	0	-		XI	5	6	1
N	V	5	15	6		XII	1	0	0
	VI	17	7	33	S	X	1	7	0
	VII	15	21	30		XI	5	29	1
O	VIII	1	0	-		XII	10	12	15
	VI	1	34	0		XIII	9	21	10
	VII	19	7	21		XIV	9	1	15
	VIII	15	30	30		XV	1	0	0
	IX	1	0	-					
total							208	409	332

TMAP can treat all elements from hydrogen to the iron group simultaneously (Rauch 1997, 2003). A small grid of model atmospheres was calculated in order to match the relevant temperature range (Fig. 1). The statistics of the model atoms which were used in the calculations are summarized in Table 1.

4. Results

The comparison of our preliminary model atmosphere fluxes with the Apr 2003 XMM-Newton RGS spectrum of V4743 Sgr (Fig. 3) with the help of XSPEC has shown that at solar abundance ratios, we cannot fit the overall flux distribution because the flux at wavelength smaller than 26 Å is too low (Figs. 2, 3). Although the fit is not perfect yet, it is obvious that the C/N abundance ratio is much smaller (about a hundred times) than the solar value, typical for CNO processed material. The fit to the observation (Fig. 3), based on our first test models (Fig. 1), is best at a T_{eff} of about 610 kK and $N_{\text{H}} = 3.5 \cdot 10^{20}/\text{cm}^{-2}$.




Figure 3. Comparison of the XMM-Newton RGS-1 spectrum with NLTE model fluxes (thick: like given in Fig. 1, thin: solar, dashed: black body). T_{eff} and N_{H} are determined with XSPEC. A black body (dashed) is shown for comparison only. The synthetic spectra are shifted by -2400 km/sec.

Acknowledgments. T.R. is supported by the DLR under grant 50 OR 0201.

References

- Haseda, K., West, D., Yamaoka, H., & Masi, G. 2002, IAU Circ. 7975
 Kato, T., Hishikura, T., West, J.D., et al. 2002, IAU Circ. 7976
 Ness, J.-U., Starrfield, S., Burwitz, V., et al. 2003, ApJ, 594, L127
 Orio, M., Leibowitz, E., Rodriguez, P., et al. 2003, IAU Circ. 8131
 Rauch, T. 1997, A&A, 320, 237
 Rauch, T. 2003, A&A, 403, 709
 Rauch, T., & Deetjen, J.L. 2003, in: Workshop on Stellar Atmosphere Modeling, eds. I. Hubeny, D. Mihalas, K. Werner, The ASP Conference Series, 288, 103
 Werner, K., Dreizler, S., Deetjen, J.L., Nagel, T., Rauch, T., & Schuh, S.L. 2003, in: Workshop on Stellar Atmosphere Modeling, eds. I. Hubeny, D. Mihalas, K. Werner, The ASP Conference Series, 288, 31